

The geometry of quantum lens spaces

Giovanni Landi

Trieste

The interrelation between MP, NT and N-CG

ESI - Vienna, March 9-13, 2015

Pimsner algebras and Gysin sequences from principal circle actions F. Arici, J. Kaad, [G.L.](#)

arXiv:1409.5335 [math.QA] ; JNcG in press

The Gysin sequence for quantum lens spaces

F. Arici, S. Brain, [G.L.](#)

arXiv:1401.6788 [math.QA] ; JNcG in press

*Anti-selfdual connections on the quantum projective plane:
Monopoles*

F. D'Andrea, [G.L.](#)

CMP 297 (2010) 841–893

Abstract:

- Quantum lens spaces as ‘direct sums of line bundles’

‘Total spaces’ of principal bundles over quantum projective spaces

- For each of these QLS a Gysin sequence in KK-theory

Used to compute the KK-theory of the QLS.s

Explicit geometric representatives of the K-theory classes which are ‘line bundles’ and generically are ‘torsion classes’

- On line bundles on QPS: monopole connections
- On higher rank bundles on QPS: instanton connections

The classical Gysin sequence

Long exact sequence in cohomology; for any sphere bundle

In particular, for circle bundles: $U(1) \rightarrow E \xrightarrow{\pi} X$

$$\dots \longrightarrow H^k(E) \xrightarrow{\pi_*} H^{k-1}(X) \xrightarrow{\cup c_1(E)} H^{k+1}(X) \xrightarrow{\pi^*} H^{k+1}(E) \longrightarrow \dots$$

complicate to generalize to quantum spaces

rather go to K-theory

Projective spaces and lens spaces

$$\mathbb{C}P^n = S^{2n+1}/U(1) \quad \text{and} \quad L^{(n,r)} = S^{2n+1}/\mathbb{Z}_r$$

assemble in principal bundles : $S^{2n+1} \longrightarrow L^{(n,r)} \xrightarrow{\pi} \mathbb{C}P^n$

This leads to the **Gysin sequence** in topological K-theory:

$$0 \longrightarrow K^1(L^{(n,r)}) \xrightarrow{\delta} K^0(\mathbb{C}P^n) \xrightarrow{\alpha} K^0(\mathbb{C}P^n) \xrightarrow{\pi^*} K^0(L^{(n,r)}) \longrightarrow 0$$

δ is a 'connecting homomorphism'

α is multiplication by the **Euler class** $\chi(\mathcal{O}_{-r}) := 1 - [\mathcal{O}_{-r}]$

From this:

$$K^1(L^{(n,r)}) \simeq \ker(\alpha) \quad \text{and} \quad K^0(L^{(n,r)}) \simeq \text{coker}(\alpha)$$

torsion groups

The quantum spheres and the projective spaces

The coordinate algebra $\mathcal{O}(S_q^{2n+1})$ of quantum **sphere** S_q^{2n+1} :
-algebra generated by $2n + 2$ elements $\{z_i, z_i^\}_{i=0, \dots, n}$ s.t.:

$$\begin{aligned} z_i z_j &= q^{-1} z_j z_i & 0 \leq i < j \leq n, \\ z_i^* z_j &= q z_j z_i^* & i \neq j, \\ [z_n^*, z_n] &= 0, \quad [z_i^*, z_i] = (1 - q^2) \sum_{j=i+1}^n z_j z_j^* & i = 0, \dots, n-1, \end{aligned}$$

and a sphere relation:

$$1 = z_0 z_0^* + z_1 z_1^* + \dots + z_n z_n^* .$$

L. Vaksman, Ya. Soibelman, 1991 ; M. Welk, 2000

The $*$ -subalgebra of $\mathcal{O}(S_q^{2n+1})$ generated by

$$p_{ij} := z_i^* z_j$$

coordinate algebra $\mathcal{O}(\mathbb{C}P_q^n)$ of the quantum **projective space** $\mathbb{C}P_q^n$

Invariant elements for the $U(1)$ -action on the algebra $\mathcal{O}(S_q^{2n+1})$:

$$(z_0, z_1, \dots, z_n) \mapsto (\lambda z_0, \lambda z_1, \dots, \lambda z_n), \quad \lambda \in U(1).$$

the fibration $S_q^{2n+1} \rightarrow \mathbb{C}P_q^n$ is a quantum $U(1)$ -principal bundle:

$$\mathcal{O}(\mathbb{C}P_q^n) = \mathcal{O}(S_q^{2n+1})^{U(1)} \hookrightarrow \mathcal{O}(S_q^{2n+1}).$$

The C^* -algebras $C(S_q^{2n+1})$ and $C(\mathbb{C}P_q^n)$ of continuous functions: completions of $\mathcal{O}(S_q^{2n+1})$ and $\mathcal{O}(\mathbb{C}P_q^n)$ in the universal C^* -norms

these are **graph algebras** **J.H. Hong, W. Szymański 2002**

$$\Rightarrow K_0(\mathbb{C}P_q^n) \simeq \mathbb{Z}^{n+1} \simeq K^0(C(\mathbb{C}P_q^n))$$

F. D'Andrea, G. L. 2010

Generators of the homology group $K^0(C(\mathbb{C}P_q^n))$ given explicitly as (classes of) even Fredholm modules

$$\mu_k = (\mathcal{O}(\mathbb{C}P_q^n), \mathcal{H}_{(k)}, \pi^{(k)}, \gamma_{(k)}, F_{(k)}), \quad \text{for } 0 \leq k \leq n.$$

Generators of the K-theory $K_0(\mathbb{C}P_q^n)$ also given explicitly as projections whose entries are polynomial functions:

line bundles & projections

For $N \in \mathbb{Z}$, vector-valued functions

$$\Psi_N := (\psi_{j_0, \dots, j_n}^N) \quad \text{s.t.} \quad \Psi_N^* \Psi_N = 1$$

$\Rightarrow P_N := \Psi_N \Psi_N^*$ is a **projection**:

$$P_N \in M_{d_N}(\mathcal{O}(\mathbb{C}P_q^n)), \quad d_N := \binom{|N| + n}{n},$$

Entries of P_N are $U(1)$ -invariant and so elements of $\mathcal{O}(\mathbb{C}P_q^n)$

Proposition 1. For all $N \in \mathbb{N}$ and for all $0 \leq k \leq n$ it holds that

$$\langle [\mu_k], [P_{-N}] \rangle := \text{Tr}_{\mathcal{H}_k}(\gamma_{(k)}(\pi^{(k)}(\text{Tr } P_{-N})) = \binom{N}{k},$$

$[\mu_0], \dots, [\mu_n]$ are generators of $K^0(C(\mathbb{C}P_q^n))$,

and $[P_0], \dots, [P_{-n}]$ are generators of $K_0(\mathbb{C}P_q^n)$

The matrix of couplings $M \in M_{n+1}(\mathbb{Z})$ is invertible over \mathbb{Z} :

$$M_{ij} := \langle [\mu_i], [P_{-j}] \rangle = \binom{j}{i}, \quad (M^{-1})_{ij} = (-1)^{i+j} \binom{j}{i}.$$

These are bases of \mathbb{Z}^{n+1} as \mathbb{Z} -modules;

they generate \mathbb{Z}^{n+1} as an Abelian group.

The inclusion $\mathcal{O}(\mathbb{C}\mathbb{P}_q^n) \hookrightarrow \mathcal{O}(S_q^{2n+1})$ is a $U(1)$ q.p.b.

To a projection P_N there corresponds an **associated line bundle**

$$\mathcal{L}_N \simeq (\mathcal{O}(\mathbb{C}\mathbb{P}_q^n))^{d_N} P_N \simeq P_{-N} (\mathcal{O}(\mathbb{C}\mathbb{P}_q^n))^{d_N}$$

\mathcal{L}_N made of elements of $\mathcal{O}(S_q^{2n+1})$ transforming under $U(1)$ as

$$\varphi_N \mapsto \varphi_N \lambda^{-N}, \quad \lambda \in U(1)$$

Each \mathcal{L}_N is indeed a bimodule over $\mathcal{L}_0 = \mathcal{O}(\mathbb{C}\mathbb{P}_q^n)$; – **the bimodule of equivariant maps** for the IRREP of $U(1)$ with **weight N** . Also,

$$\mathcal{L}_N \otimes_{\mathcal{O}(\mathbb{C}\mathbb{P}_q^n)} \mathcal{L}_M \simeq \mathcal{L}_{N+M}$$

Denote $[P_N] = [\mathcal{L}_N]$ in the group $K_0(\mathbb{C}P_q^n)$.

The module \mathcal{L}_N is a **line bundle**, in the sense that its '**rank**' (as computed by pairing with $[\mu_0]$) is equal to 1

Completely characterized by its '**first Chern number**' (as computed by pairing with the class $[\mu_1]$):

Proposition 2. *For all $N \in \mathbb{Z}$ it holds that*

$$\langle [\mu_0], [\mathcal{L}_N] \rangle = 1 \quad \text{and} \quad \langle [\mu_1], [\mathcal{L}_N] \rangle = -N.$$

The line bundle \mathcal{L}_{-1} emerges as a central character:
its only non-vanishing charges are

$$\langle [\mu_0], [\mathcal{L}_{-1}] \rangle = 1 \qquad \langle [\mu_1], [\mathcal{L}_{-1}] \rangle = 1$$

\mathcal{L}_{-1} is the *tautological line bundle* for $\mathbb{C}P_q^n$,

with *Euler class*

$$u = \chi([\mathcal{L}_{-1}]) := 1 - [\mathcal{L}_{-1}].$$

Proposition 3. *It holds that*

$$K_0(\mathbb{C}P_q^n) \simeq \mathbb{Z}[u]/u^{n+1} \simeq \mathbb{Z}^{n+1}.$$

$[\mu_k]$ and $(-u)^j$ are *dual bases* of K-homology and K-theory

The quantum lens spaces

Fix an integer $r \geq 2$ and define

$$\mathcal{O}(\mathbb{L}_q^{(n,r)}) := \bigoplus_{N \in \mathbb{Z}} \mathcal{L}_{rN}.$$

Proposition 4.

$\mathcal{O}(\mathbb{L}_q^{(n,r)})$ is a $*$ -algebra; all elements of $\mathcal{O}(S_q^{2n+1})$ invariant under the action $\alpha_r : \mathbb{Z}_r \rightarrow \text{Aut}(\mathcal{O}(S_q^{2n+1}))$ of the cyclic group \mathbb{Z}_r :

$$(z_0, z_1, \dots, z_n) \mapsto (e^{2\pi i/r} z_0, e^{2\pi i/r} z_1, \dots, e^{2\pi i/r} z_n).$$

The 'dual' $\mathbb{L}_q^{(n,r)}$:

the *quantum lens space* of dimension $2n + 1$ (and index r)

There are algebra inclusions

$$j : \mathcal{O}(\mathbb{C}P_q^n) \hookrightarrow \mathcal{O}(\mathbb{L}_q^{(n,r)}) \hookrightarrow \mathcal{O}(S_q^{2n+1}).$$

Pulling back line bundles

Proposition 5. *The algebra inclusion $j : \mathcal{O}(\mathbb{C}P_q^n) \hookrightarrow \mathcal{O}(L_q^{(n,r)})$ is a quantum principal bundle with structure group $\tilde{U}(1) := U(1)/\mathbb{Z}_r$:*

$$\mathcal{O}(\mathbb{C}P_q^n) = \mathcal{O}(L_q^{(n,r)})^{\tilde{U}(1)}.$$

Then one can ‘**pull-back**’ line bundles from $\mathbb{C}P_q^n$ to $L_q^{(n,r)}$.

$$\begin{array}{ccc} \tilde{\mathcal{L}}_N & \xleftarrow{j^*} & \mathcal{L}_N \\ \downarrow \text{dotted} & & \downarrow \text{dotted} \\ \mathcal{O}(L_q^{(n,r)}) & \xleftarrow{j} & \mathcal{O}(\mathbb{C}P_q^n). \end{array}$$

Definition 6. For each \mathcal{L}_N an $\mathcal{O}(\mathbb{C}P_q^n)$ -bimodule (a line bundle over $\mathbb{C}P_q^n$), its 'pull-back' to $L_q^{(n,r)}$ is the $\mathcal{O}(L_q^{(n,r)})$ -bimodule

$$\tilde{\mathcal{L}}_N = j_*(\mathcal{L}_N) := \mathcal{O}(L_q^{(n,r)}) \otimes_{\mathcal{O}(\mathbb{C}P_q^n)} \mathcal{L}_N.$$

The algebra inclusion $j : \mathcal{O}(\mathbb{C}P_q^n) \rightarrow \mathcal{O}(L_q^{(n,r)})$ induces a map

$$j_* : K_0(\mathbb{C}P_q^n) \rightarrow K_0(L_q^{(n,r)})$$

Each \mathcal{L}_N over $\mathbb{C}P_q^n$ is not free when $N \neq 0$,

this need not be the case for $\tilde{\mathcal{L}}_N$ over $L_q^{(n,r)}$:

the **pull-back** $\tilde{\mathcal{L}}_{-r}$ of \mathcal{L}_{-r} is **tautologically free** :

$$\tilde{\mathcal{L}}_{-r} = \mathcal{O}(L_q^{(n,r)}) \otimes_{\mathcal{L}_0} \mathcal{L}_{-r} \simeq \mathcal{O}(L_q^{(n,r)}) = \tilde{\mathcal{L}}_0.$$

$\Rightarrow (\tilde{\mathcal{L}}_{-N})^{\otimes r} \simeq \tilde{\mathcal{L}}_{-rN}$ also has trivial class for any $N \in \mathbb{Z}$

$\tilde{\mathcal{L}}_{-N}$ define **torsion classes**; they generate the group $K_0(L_q^{(n,r)})$

Multiplying by the Euler class

A second crucial ingredient

$$\alpha : K_0(\mathbb{C}P_q^n) \rightarrow K_0(\mathbb{C}P_q^n),$$

α is multiplication by $\chi(\mathcal{L}_{-r}) := 1 - [\mathcal{L}_{-r}]$

the **Euler class** of the line bundle \mathcal{L}_{-r}

Assembly these into an exact sequence, the *Gysin sequence*

$$0 \rightarrow K_1(L_q^{(n,r)}) \rightarrow K_0(\mathbb{C}P_q^n) \xrightarrow{\alpha} K_0(\mathbb{C}P_q^n) \xrightarrow{j^*} K_0(L_q^{(n,r)}) \rightarrow 0$$

$$0 \rightarrow K_1(L_q^{(n,r)}) \xrightarrow{\text{Ind}_{\mathfrak{D}}} K_0(\mathbb{C}P_q^n) \rightarrow \dots$$

and

$$\dots \rightarrow K_0(L_q^{(n,r)}) \xrightarrow{\text{Ind}_{\mathfrak{D}}} 0$$

$\text{Ind}_{\mathfrak{D}}$ comes from Kasparov theory

Write $A := C(L_q^{(n,r)})$, $F := C(\mathbb{C}P_q^n)$

The infinitesimal generator of the circle action determines an unbounded self-adjoint operator

$$\mathfrak{D} : \text{Dom}(\mathfrak{D}) \rightarrow X$$

Theorem 7. (Carey, Neshveyev, Nest, Rennie 2011)

The pair (X, \mathfrak{D}) yields a class in the bivariant $KK_1(A, F)$

the Kasparov product with the class $[(X, \mathfrak{D})]$ thus furnishes

$$\text{Ind}_{\mathfrak{D}} : K_*(A) \rightarrow K_{*+1}(F), \quad \text{Ind}_{\mathfrak{D}}(-) := -\hat{\otimes}_A [(X, \mathfrak{D})].$$

Theorem 8. (Arici, Brain, L.) *The Gysin sequence is exact*

This leads to a commutative diagram

$$\begin{array}{ccccccc}
 0 \longrightarrow & K_0(S(A)) & \xrightarrow{i_*} & K_0(M(F, A)) & \xrightarrow{ev_*} & K_0(F) & \xrightarrow{\partial} & K_1(S(A)) \longrightarrow 0 \\
 & \downarrow \text{id} & & \downarrow \simeq & & \downarrow \times[-\mathcal{L}_{-r}] & & \downarrow \text{Bott} \\
 0 \longrightarrow & K_1(A) & \xrightarrow{\text{Ind}_{\mathfrak{D}}} & K_0(F) & \xrightarrow{\alpha} & K_0(F) & \xrightarrow{j_*} & K_0(A) \longrightarrow 0
 \end{array}$$

Some practical and important applications, notably, the computation of the K-theory of the quantum lens spaces $L_q^{(n,r)}$.

Thus

$$K_1(L_q^{(n,r)}) \simeq \ker(\alpha), \quad K_0(L_q^{(n,r)}) \simeq \operatorname{coker}(\alpha).$$

Moreover, *geometric* generators of the groups

$$K_1(L_q^{(n,r)}) \quad K_0(L_q^{(n,r)})$$

for the latter as pulled-back line bundles from $\mathbb{C}P_q^n$ to $L_q^{(n,r)}$

Explicit generators as integral combinations of powers of the pull-back to the lens space $L_q^{(n,r)}$ of the generator

$$u := 1 - [\mathcal{L}_{-1}]$$

Example 9. For $n = 1$

$$K_0(C(L_q^{(1,r)})) = \mathbb{Z} \oplus \mathbb{Z}_r .$$

From definition $[\tilde{\mathcal{L}}_{-r}] = 1$, thus $\tilde{\mathcal{L}}_{-1}$ generates the torsion part.

Alternatively, from $u^2 = 0$ it follows that $\mathcal{L}_{-j} = -(j-1) + j\mathcal{L}_{-1}$; upon lifting to $L_q^{(1,r)}$, for $j = r$ this yields

$$r(1 - [\tilde{\mathcal{L}}_{-1}]) = 0$$

or $1 - [\tilde{\mathcal{L}}_{-1}]$ is cyclic of order r .

Example 10. If $r = 2$ $L_q^{(n,2)} = S_q^{2n+1}/\mathbb{Z}_2 = \mathbb{R}P_q^{2n+1}$,
the quantum real projective space, we get

$$K_0(C(\mathbb{R}P_q^{2n+1})) = \mathbb{Z} \oplus \mathbb{Z}_{2^n}$$

Owing to $\tilde{\mathcal{L}}_{-2} \simeq \tilde{\mathcal{L}}_0$ one has

$$(1 - [\tilde{\mathcal{L}}_{-1}])^2 = 2(1 - [\tilde{\mathcal{L}}_{-1}]),$$

Since $u^{n+1} = 0$, with $u = 1 - [\mathcal{L}_{-1}]$, when pulled back to the lens space, by iterating this implies that

$$0 = (1 - [\tilde{\mathcal{L}}_{-1}])^{n+1} = 2^n(1 - [\tilde{\mathcal{L}}_{-1}]);$$

the generator $1 - [\tilde{\mathcal{L}}_{-1}]$ is cyclic with the correct order 2^n .

Example 11. For $n = 2$ there are two cases.

Use $\tilde{u} = 1 - [\tilde{\mathcal{L}}_{-1}]$. Conditions $[\tilde{\mathcal{L}}_{-(r+j)}] = [\tilde{\mathcal{L}}_{-j}]$ lead to

$$\frac{1}{2}r(r-1)\tilde{u}^2 - r\tilde{u} = 0 \quad \text{and} \quad r\tilde{u}^2 = 0,$$

When $r = 2k + 1$; these say that \tilde{u} and \tilde{u}^2 are cyclic of order r :

$$r\tilde{u} = 0, \quad r\tilde{u}^2 = 0, \quad K_0(\mathcal{L}_q^{(2,r)}) = \mathbb{Z} \oplus \mathbb{Z}_r \oplus \mathbb{Z}_r$$

When $r = 2k$; $(\tilde{\mathcal{L}}_{-2})^k \simeq \tilde{\mathcal{L}}_0 \Rightarrow (1 - [\tilde{\mathcal{L}}_{-k}])^2 = 2(1 - [\tilde{\mathcal{L}}_{-k}])$, and

$$0 = (1 - [\tilde{\mathcal{L}}_{-k}])^3 = 4(1 - [\tilde{\mathcal{L}}_{-k}]) = 4k\tilde{u} - 2k(k-1)\tilde{u}^2$$

This yields $\tilde{u}^2 + 2\tilde{u}$ of order $r/2$ and \tilde{u} is of order $2r$

$$\frac{1}{2}r(\tilde{u}^2 + 2\tilde{u}) = 0, \quad 2r\tilde{u} = 0, \quad K_0(C(\mathcal{L}_q^{(2,r)})) = \mathbb{Z} \oplus \mathbb{Z}_{\frac{r}{2}} \oplus \mathbb{Z}_{2r}$$

Example 12. When $n = 3$ there are four cases

Case $r \equiv 0 \pmod{6}$:

$$K_0(C(L_q^{(3,r)})) = \mathbb{Z} \oplus \mathbb{Z}_{\frac{r}{6}} \oplus \mathbb{Z}_{\frac{r}{2}} \oplus \mathbb{Z}_{12r}$$

Case $r \equiv 2, 4 \pmod{6}$:

$$K_0(C(L_q^{(3,r)})) = \mathbb{Z} \oplus \mathbb{Z}_{\frac{r}{2}} \oplus \mathbb{Z}_{\frac{r}{2}} \oplus \mathbb{Z}_{4r}$$

Case $r \equiv 3 \pmod{6}$:

$$K_0(C(L_q^{(3,r)})) = \mathbb{Z} \oplus \mathbb{Z}_{\frac{r}{3}} \oplus \mathbb{Z}_r \oplus \mathbb{Z}_{3r}$$

Case $r \equiv 1, 5 \pmod{6}$:

$$K_0(C(L_q^{(3,r)})) = \mathbb{Z} \oplus \mathbb{Z}_r \oplus \mathbb{Z}_r \oplus \mathbb{Z}_r$$

All with explicit generators

More general scheme: Pimsner algebras M.V. Pimsner '97

The slogan: a line bundle is a self-Morita equivalence bimodule

E a (right) Hilbert module over B

B -valued hermitian structure $\langle \cdot, \cdot \rangle$ on E

$\mathcal{L}(E)$ adjointable operators; $\mathcal{K}(E) \subseteq \mathcal{L}(E)$ compact operators

with $\xi, \eta \in E$, denote $\theta_{\xi, \eta} \in \mathcal{K}(E)$ defined by $\theta_{\xi, \eta}(\zeta) = \xi \langle \eta, \zeta \rangle$

There is an isomorphism $\phi : B \rightarrow \mathcal{K}(E)$ and E is a B -bimodule

Comparing with before:

$$\mathcal{O}(\mathbb{C}P_q^n) \rightsquigarrow B \quad \text{and} \quad \mathcal{L}_{-r} \rightsquigarrow E$$

Look for the analogue of $\mathcal{O}(L_q^{(n,r)}) \rightsquigarrow \mathcal{O}_E$ Pimsner algebra

Define the B -module

$$E_\infty := \bigoplus_{N \in \mathbb{Z}} E^{\widehat{\otimes}_\phi N}, \quad E^0 = B$$

$E \widehat{\otimes}_\phi E$ the inner tensor product: a B -Hilbert module with B -valued hermitian structure

$$\langle \xi_1 \widehat{\otimes} \eta_1, \xi_2 \widehat{\otimes} \eta_2 \rangle = \langle \eta_1, \phi(\langle \xi_1, \xi_2 \rangle) \eta_2 \rangle$$

$E^{-1} = E^*$ the dual module;

its elements are written as λ_ξ for $\xi \in E$: $\lambda_\xi(\eta) = \langle \xi, \eta \rangle$

For each $\xi \in E$ a bounded adjointable operator

$$S_\xi : E_\infty \rightarrow E_\infty$$

generated by $S_\xi : E^{\widehat{\otimes}_\phi N} \rightarrow E^{\widehat{\otimes}_\phi (N+1)}$:

$$\begin{aligned} S_\xi(b) &:= \xi b, & b \in B, \\ S_\xi(\xi_1 \otimes \cdots \otimes \xi_N) &:= \xi \otimes \xi_1 \otimes \cdots \otimes \xi_N, & N > 0, \\ S_\xi(\lambda_{\xi_1} \otimes \cdots \otimes \lambda_{\xi_{-N}}) &:= \lambda_{\xi_2 \phi^{-1}(\theta_{\xi_1, \xi})} \otimes \lambda_{\xi_3} \otimes \cdots \otimes \lambda_{\xi_{-N}}, & N < 0. \end{aligned}$$

Definition 13. The *Pimsner algebra* \mathcal{O}_E of the pair (ϕ, E) is the smallest subalgebra of $\mathcal{L}(E_\infty)$ which contains the operators $S_\xi : E_\infty \rightarrow E_\infty$ for all $\xi \in E$.

Pimsner: universality of \mathcal{O}_E

There is a natural inclusion

$B \hookrightarrow \mathcal{O}_E$ a generalized principal circle bundle

roughly: as a vector space $\mathcal{O}_E \simeq E_\infty$ and

$$E^{\widehat{\otimes} \phi^N} \ni \eta \mapsto \eta \lambda^{-N}, \quad \lambda \in \mathbf{U}(1)$$

Two natural classes in KK-theory:

1. the class $[E] \in KK_0(B, B)$
of the even Kasparov module $(E, \phi, 0)$ (with trivial grading)

the map $\mathbf{1} - [E]$ has the role of the *Euler class* $\chi(E) := \mathbf{1} - [E]$

of the line bundle E over the ‘noncommutative space’ B

2. the class $[\partial] \in KK_1(\mathcal{O}_E, B)$

of the odd Kasparov module $(E_\infty, \tilde{\phi}, F)$:

$F := 2P - 1 \in \mathcal{L}(E_\infty)$ of the projection $P : E_\infty \rightarrow E_\infty$ with

$$\text{Im}(P) = \left(\bigoplus_{N=0}^{\infty} E^{\hat{\otimes}_{\phi} N} \right) \subseteq E_\infty$$

and inclusion $\tilde{\phi} : \mathcal{O}_E \rightarrow \mathcal{L}(E_\infty)$.

The Kasparov product induces group homomorphisms

$$[E] : K_*(B) \rightarrow K_*(B), \quad [E] : K^*(B) \rightarrow K^*(B)$$

and

$$[\partial] : K_*(\mathcal{O}_E) \rightarrow K_{*+1}(B), \quad [\partial] : K^*(B) \rightarrow K^{*+1}(\mathcal{O}_E),$$

Associated six-terms exact sequences **Gysin sequences**:
in K-theory:

$$\begin{array}{ccccc}
 K_0(B) & \xrightarrow{1-[E]} & K_0(B) & \xrightarrow{i_*} & K_0(\mathcal{O}_E) \\
 \uparrow [\partial] & & & & \downarrow [\partial] \\
 K_1(\mathcal{O}_E) & \xleftarrow{i_*} & K_1(B) & \xleftarrow{1-[E]} & K_1(B)
 \end{array} ;$$

the corresponding one in K-homology:

$$\begin{array}{ccccc}
 K^0(B) & \xleftarrow{1-[E]} & K^0(B) & \xleftarrow{i^*} & K^0(\mathcal{O}_E) \\
 \downarrow [\partial] & & & & \uparrow [\partial] \\
 K^1(\mathcal{O}_E) & \xrightarrow{i^*} & K^1(B) & \xrightarrow{1-[E]} & K^1(B)
 \end{array} .$$

In fact in KK-theory

Quantum weighted projective lines and lens spaces:

$B = \mathcal{O}(W_q(k, l)) =$ quantum weighted projective line

the fixed point algebra for a weighted circle action on $\mathcal{O}(S_q^3)$

$$z_0 \mapsto \lambda^k z_0, \quad z_1 \mapsto \lambda^l z_1, \quad \lambda \in \mathbf{U}(1)$$

The corresponding universal enveloping C^* -algebra $C(W_q(k, l))$ does not in fact depend on the label k : isomorphic to the unitalization of l copies of $\mathcal{K} =$ compact operators on $l^2(\mathbb{N}_0)$

$$C(W_q(k, l)) = \widetilde{\bigoplus_{s=0}^l \mathcal{K}}$$

Then: $K_0(C(W_q(k, l))) = \mathbb{Z}^{l+1}, \quad K_1(C(W_q(k, l))) = 0$

a partial resolution of singularity, since classically

$$K_0(C(W(k, l))) = \mathbb{Z}^2.$$

$\mathcal{O}_E = \mathcal{O}(L_q(lk; k, l)) = \text{quantum lens space}$

Indeed, a vector space decomposition

$$\mathcal{O}(L_q(lk; k, l)) = \bigoplus_{N \in \mathbb{Z}} \mathcal{O}_{(N)}(k, l),$$

with $E = \mathcal{O}_{(1)}(k, l)$ a right finitely projective module

$$\mathcal{O}_{(1)}(k, l) := (z_1^*)^k \cdot \mathcal{O}(W_q(k, l)) + (z_0^*)^l \cdot \mathcal{O}(W_q(k, l))$$

Also, $\mathcal{O}(L_q(lk; k, l))$ the fixes point algebra of a cyclic action

$$\mathbb{Z}/(lk)\mathbb{Z} \times S_q^3 \rightarrow S_q^3$$

$$z_0 \mapsto \exp\left(\frac{2\pi i}{l}\right) z_0, \quad z_1 \mapsto \exp\left(\frac{2\pi i}{k}\right) z_1.$$

K-theory and K-homology of quantum lens space

Denote the diagonal inclusion by $\iota : \mathbb{Z} \rightarrow \mathbb{Z}^l$, $1 \mapsto (1, \dots, 1)$ with transpose $\iota^t : \mathbb{Z}^l \rightarrow \mathbb{Z}$, $\iota^t(m_1, \dots, m_l) = m_1 + \dots + m_l$.

Theorem 14. (Arici, Kaad, L.) With $k, l \in \mathbb{N}$ coprime:

$$K_0(L_q(lk; k, l)) \simeq \text{coker}(1 - E) \simeq \mathbb{Z} \oplus (\mathbb{Z}^l / \text{Im}(\iota))$$

$$K_1(L_q(lk; k, l)) \simeq \ker(1 - E) \simeq \mathbb{Z}^l$$

as well as

$$K^0(L_q(lk; k, l)) \simeq \ker(1 - E^t) \simeq \mathbb{Z} \oplus (\ker(\iota^t))$$

$$K^1(L_q(lk; k, l)) \simeq \text{coker}(1 - E^t) \simeq \mathbb{Z}^l.$$

Again there is no dependence on the label k .

‘grand motivations / applications’ :

Gauge fields on noncommutative spaces

T-duality for noncommutative spaces

Chern-Simons theory

A Gysin sequence for $U(1)$ -bundles

relates H -flux (three-forms on the total space E) to line bundles (two-forms on the base space M) also giving an isomorphism between Dixmier-Douady classes on E and line bundles on M

Summing up:

many nice and elegant and useful geometry structures

hope you enjoyed it ; more to come soon

Thank you !!